

Calc Q2 Test 3 Review Key

1) $f(x) = 4x^3 - x^4$

$f'(x) = 12x^2 - 4x^3$

$0 = 12x^2 - 4x^3$


$0 = 4x^2(3-x)$

$x=0 \quad | \quad x=3$

$f''(x) = 24x - 12x^2$


$0 = 12x(2-x)$

$x=0 \quad | \quad x=2$



a) $(-\infty, 0) \cup (0, 3)$

b) $(3, \infty)$



c) $(0, 2)$

d) $(-\infty, 0) \cup (2, \infty)$

e) Inflection Pts

$(0, 0)$ and $(2, 16)$

$f(0) = 4(0)^3 - (0)^4$

$f(2) = 4(2)^3 - 2^4$

2) $f(x) = \frac{2x}{x-2}$

$f'(x) = \frac{(x-2)(2) - (2x)(1)}{(x-2)^2}$

$f''(x) = \frac{(x-2)^2(0) - (-4)(2(x-2)(1))}{(x-2)^4}$

$f'(x) = \frac{2x-4-2x}{(x-2)^2}$

$f''(x) = \frac{0 + 8(x-2)}{(x-2)^4}$

$f'(x) = \frac{-4}{(x-2)^2}$


$f''(x) = \frac{8(x-2)}{(x-2)^4} = \frac{8}{(x-2)^3}$

$f'(x) = 0$
Never

$f'(x) = \text{und.}$
 $x-2=0$
 $x=2$


$f''(x) = 0$
Never

$f''(x) = \text{und.}$
 $x-2=0$
 $x=2$



a) Never

b) $(-\infty, 2) \cup (2, \infty)$



c) $(2, \infty)$

d) $(-\infty, 2)$

e) Inflection pt at $x=2$
But function DNE there



$$3) f(x) = (2x-5)^3 \quad f'(x) = 3(2x-5)^2(2) \quad f''(x) = 12(2x-5)(2)$$

$$f'(x) = 6(2x-5)^2 \quad f''(x) = 24(2x-5)$$

$$0 = 6(2x-5)^2 \quad 0 = 24(2x-5)$$

$$0 = 2x-5 \quad | \quad x = \frac{5}{2}$$

$$\frac{5}{2} = x$$



a) $(-\infty, \frac{5}{2}) \cup (\frac{5}{2}, \infty)$

b) Never

c) $(\frac{5}{2}, \infty)$

d) $(-\infty, \frac{5}{2})$

e) Inflection P+ @ $x = \frac{5}{2}$

$$f(\frac{5}{2}) = (2(\frac{5}{2}) - 5)^3$$

$$f(\frac{5}{2}) = 0$$

$$(\frac{5}{2}, 0)$$

$$4) f(x) = \frac{x^2}{x^2-4} \quad f'(x) = \frac{(x^2-4)(2x) - (x^2)(2x)}{(x^2-4)^2} \quad f''(x) = \frac{(x^2-4)^2(-8) - (-8x)(2)(x^2-4)(2x)}{(x^2-4)^4}$$

$$f'(x) = \frac{2x^3 - 8x - 2x^3}{(x^2-4)^2}$$

$$f''(x) = \frac{-8(x^2-4)^2 + 32x^2(x^2-4)}{(x^2-4)^4}$$

$$f'(x) = \frac{-8x}{(x^2-4)^2}$$


$$f''(x) = \frac{-8(x^2-4) + 32x^2}{(x^2-4)^3}$$

$$f'(x) = 0 \quad f'(x) = \text{und}$$

$$-8x = 0 \quad x^2 - 4 = 0$$

$$x = 0 \quad x = \pm 2$$

$$f''(x) = \frac{-8x^2 + 32 + 32x^2}{(x^2-4)^3} = \frac{24x^2 + 32}{(x^2-4)^3}$$



$$f''(x) = 0$$

$$24x^2 + 32 = 0$$


$$8(3x^2 + 4) = 0$$

$$x^2 = -\frac{4}{3} \text{ Never}$$

$$f''(x) = \text{und}$$

$$x^2 - 4 = 0$$

$$x = \pm 2$$



a) $(-\infty, -2) \cup (-2, 0)$

b) $(0, 2) \cup (2, \infty)$

c) $(-\infty, -2) \cup (2, \infty)$ d) $(-2, 2)$ e) None

5) $f(x)$ is increasing when $f'(x)$ is positive (above the x-axis)
 $(-1, 3) \cup (5, \infty)$

6) $f(x)$ is decreasing when $f'(x)$ is negative (below the x-axis)
 $(-\infty, -1) \cup (3, 5)$

b) $f(x)$ is concave up when $f'(x)$ is increasing
 $(-\infty, \frac{1}{2}) \cup (4, \infty)$

$f(x)$ is concave down when $f'(x)$ is decreasing
 $(\frac{1}{2}, 4)$

c) $f(x)$ has an inflection pt when $f'(x)$ has a max or min
Inflection pt when $x = \frac{1}{2}$ and $x = 4$

7) I \rightarrow C

II \rightarrow B

III \rightarrow A

8) a) $f(x) = 2x^3 - 9x^2 + 12x - 5$ $[0, 2]$ $f(0) = 2(0)^3 - 9(0)^2 + 12(0) - 5 = (-5)$ Abs Min
 $f'(x) = 6x^2 - 18x + 12$ $f(1) = 2(1)^3 - 9(1)^2 + 12(1) - 5 = (0)$ Abs Max
 $0 = 6(x^2 - 3x + 2)$ $f(2) = 2(2)^3 - 9(2)^2 + 12(2) - 5 = -1$
 $(x-2)(x-1)$
 $x=2 \quad x=1$

b) $f(x) = 2x^3 - 15x^2 + 24x + 2$ $[0, 2]$ $f(0) = 2(0)^3 - 15(0)^2 + 24(0) + 2 = (2)$ Abs Min
 $f'(x) = 6x^2 - 30x + 24$ $f(1) = 2(1)^3 - 15(1)^2 + 24(1) + 2 = (13)$ Abs Max
 $6(x^2 - 5x + 4)$ $f(2) = 2(2)^3 - 15(2)^2 + 24(2) + 2 = 6$
 $(x-4)(x-1)$
 $x=4 \quad x=1$

Not in
interval

9) a) $f'(-2) < 0$

b) $f'(-1) > 0$

c) $f'(0) = 0$

d) $f(1.5) > 0$

e) $f'(-3) > f'(-1)$ Tangent line @ $f'(-3)$ is steeper than @ $f'(-1)$

f) $f(1) > f'(1)$ $f(1) = 0$

g) $f''(1) > f'(1)$ $f(x)$ is concave up at $x=1$
therefore $f''(x)$ is positive

h) $f'(1) < f'(0)$